



# Assessment of composted kitchen waste and poultry manure amendments on growth, yield and heavy metal uptake by Jute mallow *Corchorus olitorius* Linn.

Oladele A. Oguntade<sup>1</sup> · Titilola S. Olagbenro<sup>1</sup> · Olubunmi A. Odusanya<sup>1</sup> · Solomon O. Olagunju<sup>1</sup> · Kayode M. Adewusi<sup>1</sup> · Adedotun T. Adegoke<sup>1</sup>

Received: 31 December 2017 / Accepted: 12 November 2018 / Published online: 17 November 2018

© The Author(s) 2018

## Abstract

**Purpose** This study was carried out to compare the effects of composted kitchen waste (KW) and poultry manure (PM) soil amendments on growth, yield and heavy metal uptake in edible leaf of *Corchorus olitorius*.

**Methods** Kitchen waste and PM composted for 8 weeks were applied as soil amendment at the rate of 0 (no amendment), 5, 10 and 15 t ha<sup>-1</sup>. *Corchorus* seeds were sown 2 weeks after incorporation of amendment. Compost amendments were the main treatments while application rates were in sub-plots. Treatments were replicated three times in a randomized complete block design. *Corchorus* were harvested 6 weeks after sowing, following which growth and yield parameters were measured. Concentrations of manganese (Mn), iron (Fe), copper (Cu) and zinc (Zn) in leaf tissue and plant uptake were also determined.

**Results** Result showed that PM increased *Corchorus* leaf production than KW. Poultry manure significantly ( $p \leq 0.01$ ) increased Cu accumulation in leaf. Bioaccumulation of Cu was 0.005 and 0.011 mg kg<sup>-1</sup> for KW and PM, respectively. The metals except Zn in leaf tissue showed significant ( $p \leq 0.01$ ) correlation with both fresh and dry weights of *Corchorus*. Despite high concentration of heavy metals in the compost, bioaccumulation in leaf was lower than maximum allowable limit of 0.1 mg kg<sup>-1</sup> for Cu and 0.3 mg kg<sup>-1</sup> for Mn, Fe and Zn in vegetables by WHO/FAO/FEPA.

**Conclusion** Soil amended with composted KW and PM promoted *Corchorus* growth and yield. Accumulations of heavy metals in leaf tissue are within allowable limit for vegetables.

**Keywords** Bioaccumulation · Compost · *Corchorus olitorius* · Edible leaf · Soil amendment

## Introduction

Management of domestic solid waste generated in most Nigerian homes is a growing challenge that requires urgent attention. Poor management of domestic waste impacts on our environment negatively. With larger proportion of the domestic wastes being organic, an alternative source of compostable materials that could serve as soil ameliorant could become attractive. Use of compostable solid waste as soil amendment in crop production is a sustainable way of recycling that reduces disposal to landfill (Petersen et al. 2003;

Chen and Jiang 2014). Compost is an important resource for soil amendment in organic farming with the benefits of being environment-friendly. Composted manure helps to modify physical, chemical, and biological properties of the soil, provides slow release of nutrients and increases crop yield (Agbede et al. 2008; Moral et al. 2009). Composted organic manure especially from poultry wastes have been reported to improve mineral composition in tissues of vegetables such as Jute mallow (Mazen et al. 2010; Jonathan et al. 2012). In contrast to the beneficial roles of compost, it could also be a potential source of contaminants depending on its origin (Petersen et al. 2003; Hanč et al. 2008).

Leafy vegetables have the potentials of accumulating heavy metals in their edible and non-edible portions in concentrations that could cause health problems to consumers when grown in contaminated medium (Alam et al. 2003; Oguntade et al. 2015). Uptake of heavy metals by vegetables in high concentration could also hinder metabolic processes

✉ Oladele A. Oguntade  
deletade@yahoo.com; dele.oguntade@oouagoiwoye.edu.ng

<sup>1</sup> Department of Crop Production, College of Agricultural Sciences, Olabisi Onabanjo University, P.M.B. 0012, Ayetoro Campus, Ayetoro, Ogun State, Nigeria

such as photosynthesis and growth in such plant and consequently lead to reduction in yield (Yang et al. 2002; Wang et al. 2013; Ng et al. 2016). *Corchorus olitorius* known as Jute mallow is a popular green leafy vegetable consumed by various communities in many African countries including Nigeria (Schippers 2002; Samra et al. 2007). The slimy sticky sauce of its cooked leaves is often considered as the best and more acceptable one than okro in western Nigeria even at social gathering for consumption of any starchy staples served (Denloye et al. 2014). *Corchorus olitorius* is a vegetable consumed for its dietary fiber, vitamins, minerals, protein as well as abundant  $\beta$ -carotene and other carotenoids needed for good health (Schippers 2002). The leaf is also a rich source of iron, protein, calcium, thiamin, riboflavin, niacin, foliate, and dietary fiber (Oyedele et al. 2006; Samra et al. 2007).

Considering the economic and environmental benefits of organic wastes, the use of composted organic waste in production of *Corchorus olitorius* could be an alternative strategy of waste management and disposal. Although, information on the use of manure for vegetable production is well documented, little is, however, known on the use of composted kitchen waste and poultry manure for jute mallow production. This study, therefore, focuses on the growth, yield and heavy metal bioaccumulation in edible leaf of *Corchorus olitorius* grown on soil amended with composted kitchen waste and poultry manure.

## Materials and methods

### Description of the study site

This study was conducted at the Teaching and Research Farm of the Olabisi Onabanjo University, College of Agricultural Sciences, Ayetoro Campus, Ogun State, Nigeria. Ayetoro is in the derived savannah ecology of Ogun State with mean annual rainfall of 1250 mm and mean temperature of 26 °C. It has a bimodal rainfall pattern with its peaks in June and September. The experimental site was located at 07.23076°N, 03.04540°E and 89 m above sea level.

### Collection and preparation of kitchen waste and poultry manure as compost

Decomposable kitchen wastes collected from various homes and from leftovers of food at social gatherings were used for this study. Similarly, poultry manure from birds fed with maize and groundnut cake-based feed was collected from the University's poultry farm at Ayetoro campus for the compost preparation. The kitchen waste and poultry manure were composted for 8 weeks following the Chinese rural composting-high temperature method (Misra et al. 2003).

Composts were then air-dried for 2 weeks and sieved with 2 mm sieve to remove undecomposed materials before field application (Plate 1). Sub-samples of the composted wastes were analyzed in the laboratory for its chemical properties (Table 1).

### Plot establishment, soil sample collection and preparation

The experiment was a split-plot arrangement in a Randomized Complete Block Design (RCBD) replicated three times. Main-plot treatments were the composted kitchen waste and poultry manure, while sub-plot treatments were compost rates of 0 (no amendment/control), 5, 10, 15 t ha<sup>-1</sup>. The treatment plots were laid out with a spacing of 1 m between treatments and 2 m between replicates. Each of the treatments (compost) was incorporated into the soil on 1 m by 1 m elevated bed 2 weeks before sowing (WBS). Prior to treatments application, core soil samples were randomly collected (0–15 cm) from the experimental plot with a soil auger and bulked to obtain composite sample. Thereafter, the composite soil sample was air-dried and sieved with 2 mm sieve for routine analysis in the laboratory.

### Corchorus seeds preparation and sowing

Prior to sowing, *Corchorus* seeds were pre-treated with hot water at 90 °C for 4 s to break its dormancy and later air-dried for a day (Jayasuriya et al. 2007; Tolorunse et al. 2015). The air-dried seeds were then drilled on elevated beds earlier amended with either composted KW or PM at the spacing of about 10 cm between rows. Due to irregular rainfall between the second and the fourth week after sowing

**Table 1** Chemical properties of composted kitchen waste and poultry manure

Chemical composition	Units	Kitchen waste	Poultry manure
pH (H <sub>2</sub> O)		8.80	8.75
Total N	g kg <sup>-1</sup>	2.90	3.20
Total organic carbon		50.40	55.90
Total P	mg kg <sup>-1</sup>	0.46	0.35
C/N ratio		17.38	17.47
Ca	cmol kg <sup>-1</sup>	2.83	3.05
Mg		0.74	0.57
Na		1.51	0.71
K		1.46	0.69
Heavy metals			
Mn	mg kg <sup>-1</sup>	4000.00	7000.00
Fe		4200.00	6800.00
Cu		30.00	45.00
Zn		278.00	354.00

(WAS), each treatment plot was irrigated with equal volume (10 L) of fresh water collected from adjoining stream every evening. Overhead method of irrigation was adopted with the use of watering can. This is similar to the practice of local farmers in the area. Corchorus seedlings were thinned at 2 WAS to 1 plant per stand at about 4 cm within the row. Weeding was done manually by uprooting weeds regularly and retaining them on their various plots.

### Agronomic data collection

At 6 WAS, agronomic parameters of twenty (20) stands of *Corchorus olitorius* taken from a quadrant of 0.5 m by 0.5 m placed on each treatment plot were recorded. Plant height was measured with a meter rule from the soil surface to the terminal bud. The number of leaves and primary branches was determined by direct counting. The edible portions of the leaves excluding the petiole were then plucked and the fresh weight was determined. Thereafter, the leaves were cleaned with fresh tap water, air-dried and later oven-dried at 60 °C to constant weight to determine the dry weight. After drying, leaves were ground with pestle and mortar for chemical analysis.

### Chemical analysis of compost manure

The pH of the composted manure was determined with the glass electrode pH meter. Nitrogen (N) was determined by Kjeldahl method, organic carbon by wet oxidation method (Nelson and Sommer 1996), while phosphorus (P) was determined colourimetrically. The calcium (Ca), magnesium (Mg), potassium (K) and sodium (Na), as well as Mn, Fe, Cu, and Zn in the compost samples were determined by atomic absorption spectrophotometer, AAS (Buck Scientific 210VGP Model, Inc., East Norwalk, CT, USA) according to Udo et al. (2009).

### Soil analysis

The pH of the soil in water was determined by glass electrode pH meter. Total N was determined by micro-Kjeldahl method, organic carbon content was determined by Walkley and Black method (1934) using dichromate ( $K_2Cr_2O_7$ ) as oxidizing agent. Available P was determined colourimetrically by Bray-1- method (Bray and Kurtz 1945). Exchangeable bases ( $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$  and  $Na^+$ ) were extracted with 1 N ammonium acetate ( $NH_4OAc$  pH 7.0) after which  $Ca^{2+}$  and  $Mg^{2+}$  were read on AAS, while  $K^+$  and  $Na^+$  were determined with flame photometer. Exchangeable acidity was determined by titration method following extraction of soil samples with 1 N KCl solution (Anderson and Ingram 1993). The effective cation exchange capacity (ECEC) was estimated by summation of exchangeable acidity ( $H^+ + Al^{3+}$ )

and exchangeable bases ( $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$  and  $Na^+$ ). Base saturation was calculated as the ratio of sum of exchangeable cations to ECEC expressed in percentage. The heavy metals including Fe, Cu, Zn, and Mn in the soil were determined by wet digestion method and read on AAS. The analyses followed the procedures in Okalebo et al. (1993) and Udo et al. (2009). Particle size distribution was analyzed by the hydrometer method (Bouyoucos 1951).

### Heavy metal analysis in plant samples

Earlier ground Corchorus leaves were weighed (0.5 g) into digestion tubes; then 10 ml of mixture of nitric and perchloric acid (2:1) was added. The content in the tubes was digested for the first 90 min at 150 °C. Thereafter, 2 ml of concentrated HCl and distilled water was added and the temperature of the digest was raised to 230 °C and further digested for another 30 min. The digest was then allowed to cool at room temperature after which the content was transferred into a 50-ml volumetric flask and made up to mark with distilled water. The concentrations of Mn, Fe, Cu and Zn in the sample solutions were then determined using atomic absorption spectrophotometer. Plant uptake was estimated on dry weight basis as the product of metal concentration and dry weight.

### Data analysis

The data obtained were subjected to analysis of variance (ANOVA). Treatment means were separated using Fisher's Protected Least Significant Difference (LSD) at 5% probability.

## Results

### Chemical composition of compost manure

Chemical compositions of the composted KW and PM are presented in Table 1. The pH of the two composts was moderately alkaline with PM being more alkaline than KW. The total N was 2.90 g  $kg^{-1}$  in KW and 3.20 g  $kg^{-1}$  in PM. Organic carbon was lower in KW than in PM. Total phosphorus was higher in KW than in PM. Poultry manure and KW composts had the same C/N ratio level, which reflected its carbon and N contents. Except for Ca, the total Mg, Na and K were found to be higher in the KW than in PM. The concentrations of heavy metals including Mn, Fe, Cu and Zn of 4000, 4200, 30 and 278 mg  $kg^{-1}$ , respectively, in KW were lower than 7000, 6800, 45 and 354 mg  $kg^{-1}$ , respectively, in PM (Table 1).

Physicochemical soil properties of the experimental site indicated that pH of the soil was moderately acidic

**Table 2** Physicochemical soil properties of the experimental site

Soil properties	Units	Values
pH (H <sub>2</sub> O)		5.95
Total N	g kg <sup>-1</sup>	0.90
Total organic carbon		11.79
Available P	mg kg <sup>-1</sup>	17.68
Ca	cmol kg <sup>-1</sup>	8.38
Mg		2.21
Na		0.50
K		0.30
Exchangeable acidity		0.08
ECEC		11.47
Base saturation	%	99.93
Mn	mg kg <sup>-1</sup>	16.80
Fe		20.80
Cu		0.70
Zn		4.80
Sand	g kg <sup>-1</sup>	828.00
Silt		54.00
Clay		118.00
Texture		Loamy sand

(Table 2). Total N and organic carbon were very low, while available P was high. The exchangeable cations in the soil which were 8.38, 2.21, 0.50 and 0.30 cmol kg<sup>-1</sup> for Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup> and K<sup>+</sup>, respectively, were high. The exchangeable acidity was very low. Available form of Mn in the soil was 16.80 mg kg<sup>-1</sup>. The available Fe and Zn of 20.80 and 4.80 mg kg<sup>-1</sup>, respectively, in the soil were quite high. The

texture of the soil was loamy sand with high proportion of sand, low silt and moderately low clay content (Table 2).

Table 3 shows the effects of different rates of composted KW and PM amendments on the growth and yield of *Corchorus olitorius*. The effects of the two amendments on growth and yield parameters were not significantly different except on the number of leaves. Poultry manure significantly ( $p \leq 0.05$ ) increased the number of leaves in *Corchorus* compared to KW. The different rate of amendment applied increased plant height, number of leaves and primary branches significantly ( $p \leq 0.01$ ) (Table 3). The *Corchorus* grown in soil without amendment gave the least mean plant height, number of leaves and primary branches, while those grown with 15 t ha<sup>-1</sup> of the amendment gave the highest growth parameters recorded. Meanwhile, there were no significant differences on both fresh and dry biomass yields at the different rate of amendment.

The interactive effects of the type of compost and its rates of application contributed significantly ( $p \leq 0.01$ ) to increased *Corchorus* height and number of primary branches (Figs. 1 and 2). In particular, at compost rate of 15 t ha<sup>-1</sup>, the mean plant height was higher with PM than with KW (Fig. 1). Meanwhile, the difference in mean number of primary branches observed at 10 t ha<sup>-1</sup> of the amendment showed that PM was higher than KW in inducing primary branches in *Corchorus* (Fig. 2).


Accumulation of heavy metals in leaf tissue (dry weight) of *Corchorus* grown on compost amended soil is presented in Table 4. Among the metals, only Cu was significantly ( $p \leq 0.01$ ) taken up and bioaccumulated in leaf tissue of *Corchorus*. The amount of Cu which accumulated in the leaf of *Corchorus* grown on PM amended soil was

**Table 3** Effects of compost types and their rates of application on growth and yield of *Corchorus olitorius* at 6WAS


Source of variation	Plant height (cm)	Number of leaves	Number of primary branches	Leaf fresh weight (t ha <sup>-1</sup> )	Leaf dry weight (t ha <sup>-1</sup> )
Compost type					
KW	20.61	7.76	3.35	1.67	0.54
PM	20.96	8.75	3.36	1.72	0.55
(Sig.)	ns	*	ns	ns	ns
LSD <sub>(0.05)</sub>	1.01	0.78	0.08	0.60	0.17
Rates					
0	17.55	7.18	2.36	1.43	0.47
5	19.53	7.88	3.19	1.62	0.53
10	21.08	8.42	3.60	2.06	0.63
15	24.96	9.54	4.28	1.68	0.54
(Sig.)	**	**	**	ns	ns
LSD <sub>(0.05)</sub>	1.42	1.10	0.11	0.85	0.23
Compost * rate	**	ns	**	ns	ns

ns not significant

\*\*\*Significant at 5% and 1%, respectively, using Fisher's protected LSD



**Fig. 1** Responses of *Corchorus* height to rates of compost amendment. Vertical bars stand for LSD at  $p \leq 0.05$



**Fig. 2** Responses of primary branches of *Corchorus* to rates of compost amendment. Vertical bars stand for LSD at  $p \leq 0.05$

more than double compared to  $0.005 \text{ mg kg}^{-1}$  with KW amendment. The results also showed that various rates of compost amendment used were not significantly different on the heavy metal uptake in *Corchorus* leaf. However, interactive effects of the compost types and its rates further showed significant ( $p \leq 0.05$ ) uptake and accumulation of Cu in *Corchorus* leaf. The study showed that positive and significant ( $p \leq 0.01$ ) correlation exists between the heavy metals uptake in leaf tissue and biomass yield of *Corchorus* (Table 5). Both fresh and dry biomass of *Corchorus* leaf were highly correlated ( $p \leq 0.01$ ) with the heavy metals uptake except for Zn. Manganese and Zn uptake correlated significantly ( $p \leq 0.05$ ) with the number


of primary branches, while Cu and Mn uptake were correlated ( $p \leq 0.05$ ) with the plant height. Only Cu also showed significant correlation with mean number of leaves produced by *Corchorus*.

## Discussion

The proximate composition of composted KW and PM applied as soil amendment in this study revealed that PM was higher in its organic constituent than KW. This explains why PM promoted significant increase in the number of leaves in *Corchorus* compared to KW. The higher concentration of total P, Mg, Na and K found in the composted KW relative to PM was an indication of the presence of salts of these nutrients in the KW than PM. This showed that KW can serve as a good alternative to PM in supplying mineral nutrients required for growth and structural development of crops. This probably explains why there was no significant difference between KW and PM on growth and yield recorded except on the number of leaves. However, higher concentration of heavy metals in PM than KW observed could be primarily from the feeds and drugs given to the poultry birds (Zhang et al. 2005; van Zanten et al. 2014). High concentration of heavy metals in PM compared to KW could, therefore, limit its use to reduce bioaccumulation of the metals in edible portion of *Corchorus*. Since heavy metals are known to be resistant to biodegradation with long biological half-life, coupled with the tendency of been taken up by *Corchorus olitorius* from amended soil, it follows that the metals can persist in the growth medium and contaminate the food chain (Garbarino et al. 2003; Jackson et al. 2003; Sapkota et al. 2007).

The poor quality of the soil used for this study, as depicted by low concentration of nitrogen when compared with critical level of  $1.5\text{--}2.0 \text{ g kg}^{-1}$  reported by Sobulo and Osiname (1981), justifies the need for organic amendment in such soil for production of leafy vegetable like *Corchorus*. Application of composted KW and PM as soil amendments increased *Corchorus* growth and yield in comparison with the control soil without amendment. However, the high ECEC of the soil indicated that the soil exchange site was dominated by exchangeable bases particularly  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . Calcium,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  and  $\text{K}^+$  concentrations in the soil were higher than critical levels of  $1.5 \text{ cmol kg}^{-1}$  for  $\text{Ca}^{2+}$  (Enwezor et al. 1990),  $0.28 \text{ cmol kg}^{-1}$  for  $\text{Mg}^{2+}$  (Agboola and Ayodele 1985; FMANR 1990),  $0.02 \text{ cmol kg}^{-1}$  for  $\text{Na}^+$  (Amalu 1997) and  $0.15 \text{ cmol kg}^{-1}$  for  $\text{K}^+$  (Uponi and Adeoye 2000). Meanwhile, the concentration of Mn was higher than critical level of  $1.00 \text{ mg kg}^{-1}$  (Udoh et al. 2008). Iron was in several folds higher than critical level of  $4.50 \text{ mg kg}^{-1}$  reported by Lindsay and Norvell (1978). Available Cu in

**Plate 1** Composted kitchen waste (a) and poultry manure (b) used as soil amendment



**Table 4** Heavy metal uptake by *Corchorus olitorius* leaves grown on soil amended with composted kitchen waste and poultry manure 6WAS

Source of variation	Mn (mg kg <sup>-1</sup> )	Fe (mg kg <sup>-1</sup> )	Cu (mg kg <sup>-1</sup> )	Zn (mg kg <sup>-1</sup> )
Compost type				
Kitchen waste	0.014	0.210	0.005	0.016
Poultry manure	0.015	0.212	0.011	0.010
(Sig.)	ns	ns	**	ns
LSD <sub>(0.05)</sub>	0.005	0.067	0.003	0.015
Rates (t ha <sup>-1</sup> )				
0	0.011	0.178	0.006	0.006
5	0.013	0.196	0.006	0.008
10	0.018	0.254	0.010	0.012
15	0.017	0.218	0.009	0.025
(Sig.)	ns	ns	ns	ns
LSD <sub>(0.05)</sub>	0.007	0.094	0.004	0.022
Compost * rates				
	ns	ns	*	ns
FAO/WHO/FEPA maximum limits	0.30	0.30	0.10	0.30

ns not significant, FEPA Federal Environmental Protection Agency, Nigeria

\*\*\* Significant at 5% and 1%, respectively, using Fisher's protected LSD

**Table 5** Correlation analysis between yield parameters and heavy metal uptake in *Corchorus olitorius* leaves

Parameters	Cu	Fe	Mn	Zn	LDW (t ha <sup>-1</sup> )	LFW (t ha <sup>-1</sup> )	NPB	PHt. (cm)	NL
Cu									
Fe	0.71**	–							
Mn	0.75**	0.97**	–						
Zn	–0.02ns	0.15ns	0.24ns	–					
LDW (t ha <sup>-1</sup> )	0.68**	0.98**	0.94**	0.12ns	–				
LFW (t ha <sup>-1</sup> )	0.71**	0.98**	0.95**	0.10ns	0.99**	–			
NPB	0.33ns	0.25ns	0.43*	0.44*	0.18ns	0.20ns	–		
PHt. (cm)	0.46*	0.34ns	0.47*	0.11ns	0.27ns	0.28ns	0.78**	–	
NL	0.47*	0.16ns	0.31ns	0.30ns	0.12ns	0.31ns	0.73**	0.63**	–

LDW leave dry weight, LFW leaves fresh weight, NPB number of primary branches, PHt. plant height, NL number of leaves, ns not significant

\*\*\* Correlation is significant at 5% and 1%, respectively, using Fisher's protected LSD

this soil was above the critical value of  $0.2 \text{ mg kg}^{-1}$  given by Sims and Johnson (1991). Similarly, concentration of Zn in the soil was higher than critical value of  $0.8 \text{ mg kg}^{-1}$  reported by Lindsay and Norvell (1978) and  $0.5 \text{ mg kg}^{-1}$  according to Udoh et al. (2008).

The contribution of compost amendment to higher growth and yield of Corchorus compared to control soil without amendment was a reflection of the beneficial roles played by the compost amendment in enhancing crop yield (Chiu et al. 2006; Agbede et al. 2008). Although, beyond  $10 \text{ t ha}^{-1}$  rate, PM-amended soil contributed to higher yield of Corchorus than KW, it may not be safe to use higher rate of PM as soil amendment since PM contains higher concentration of heavy metals. This, in a way, can reduce the source of contaminant and subsequent transfer of heavy metal to edible portion of Corchorus. Previous studies have reported that the origin of a particular compost material could be a potential source of heavy metals in the plant tissues (Petersen et al. 2003; Hanč et al. 2008).

The higher accumulation of Cu in Corchorus leaf with PM than KW reflected the higher concentration of Cu in PM amendment over KW. This observation implies that larger proportion of Cu from PM amended soil was translocated through roots to the leaf probably due to its high mobility. This finding is contrary to what Hanč et al. (2008) found in biomass of oat grown on soil amended with compost having high level of Cu. A lower uptake of Cu reported was adduced to localization in root tissue of oat. This showed that Cu uptake and mobility to upper part of the plant are not consistent and could vary with the type of crop. The declined Mn, Fe and Cu uptake at higher compost rate of  $15 \text{ t ha}^{-1}$ , despite higher concentration of the metals in the soil, corroborates previous studies on the chelating impact of compost and organic amendments on heavy metal uptake by crops (Angelova et al. 2010). Although, the compost used in this study, especially the PM, had high levels of heavy metals, bioaccumulation in Corchorus leaf was lower than the maximum allowable limits by WHO/FAO/FEPA in vegetables. This was due to chelating reaction of the organic compounds in the compost or the role played by humic portion of the compost which immobilized the heavy metals (Tlustoš et al. 2006; Cooper et al. 2011; Oguntade et al. 2015). This showed that production of Corchorus with these composts under this condition was safe for human consumption without threat to health of the consumers.

The significant correlation of biomass yield with heavy metal uptake by Corchorus indicated that Mn, Fe and Cu contributed to leaf development and yield. In a similar study (Hanč et al. 2008), an increased in Cu uptake was found to be in close correlation with biomass yield of oat. Moreover, the correlation of Mn and Zn with the number

of primary branches indicated that these metals induced primary branching in Corchorus.

## Conclusion

It is evident in this study that composted KW can support Corchorus production favourably like composted PM. The higher concentration of P, K, Mg and Na in composted KW than PM in this study is worthy of note. Although, composted PM had higher concentration of nitrogen and calcium than KW, the higher concentrations of heavy metals in the PM could be a major source of contaminant. Hence, usage of PM in vegetable production could contribute largely to heavy metal contamination of food crops than KW. Application of  $10 \text{ t ha}^{-1}$  of composted KW and PM is suggested as soil amendment for optimum yield of Corchorus to reduce soil and food chain contamination by the toxic metals. However, while converting compostable wastes into organic fertilizer, it is imperative to investigate the potential heavy metal status of any waste to be used in production of food crops to reduce possible transfer of toxic metals to human.

**Acknowledgements** The authors of this paper are thankful to the Laboratory Technologist at the Crop/Soil Science Laboratory, College of Agricultural Sciences, Ayetoro Campus. We also appreciate Technologist at SMO Laboratory Services, Ibadan. Authors also acknowledge the efforts of Dr. AL Nassir for the editorial work on the manuscript.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Agbede TM, Ojeniyi SO, Adeyemo AJ (2008) Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in South-West, Nigeria. *Am Eurasian J Sust Agric* 2:72–77
- Agboola AA, Ayodele OJ (1985) Prospects and problem of using soil testing for adoption of fertilizer use in Ekiti Akoko. *Agricultural Development Project area*. In: Ohun HW, Nagy JG (eds) *Proceedings of workshop on appropriate technologies for farmers in semi-arid West Africa*. Purdue University Press, West Lafayette, pp 123–136
- Alam MGM, Snow ET, Tanaka A (2003) Arsenic and heavy metal contamination of vegetables grown in Samta village, Bangladesh. *Sci Total Environ* 308:83–96. [https://doi.org/10.1016/S0048-9697\(02\)00651-4](https://doi.org/10.1016/S0048-9697(02)00651-4)
- Amalu UC (1997) Evaluation of properties of selected soils of Cross River State area and their management for increased cassava yields. *Glob J Pure Appl Sci* 4(3):243–249
- Anderson JM, Ingram JSI (1993) Soil organic matter and organic carbon. In: Anderson JM, Ingram JSI (eds) *Tropical soil biology and*



- fertility: a handbook of method of analysis. CAB International, Wallingford, UK, pp 62–70
- Angelova V, Ivanova R, Pevicharova G, Ivanov K (2010) Effect of organic amendments on heavy metal uptake by potato plants. In: Proceedings of the 19th world congress of soil science, soil solution for a change world, 1–6 August, Brisbane, Australia, Published on DVD, pp 84–87
- Bouyoucos GH (1951) A recalibration of the hydrometer for testing mechanical analysis of soil. *Agron J* 43:434–438. <https://doi.org/10.2134/agronj1951.00021962004300090005x>
- Bray R, Kurtz L (1945) Determination of total organic and available forms of phosphorus in soils. *Soil Sci* 59:39–45. <https://doi.org/10.1097/00010694-194501000-00006>
- Chen Z, Jiang X (2014) Microbiological safety of chicken litter or chicken litter-based organic fertilizers: a review. *Agriculture* 4(1):1–29. <https://doi.org/10.3390/agriculture4010001>
- Chiu KK, Ye ZH, Wong MH (2006) Growth of *Vetiveria zizanioides* and *Phragmites australis* on Pb/Zn and Cu mine tailing amended with manure compost and sewage sludge: a greenhouse study. *Bioresour Technol* 97:158–170. <https://doi.org/10.1016/j.biortech.2005.01.038>
- Cooper J, Sanderson R, Cakmak I, Ozturk L, Shotton P, Carmichael A, Sadrabadi R, Tetard-Jones C, Volakakis N, Eyre M, Leifert C (2011) Effects of organic and conventional crop rotation, fertilization and crop protection practices on metal contents in wheat *Triticum aestivum*. *J Agric Food Chem* 59:4715–4724. <https://doi.org/10.1021/jf104389m>
- Denloye AA, Makinde OSC, Ajelara KO, Alafia AO, Oiku EA, Dosumu OA, Makanjuola WA, Olowu RA (2014) Insects infesting selected vegetables in Lagos and the control of infestation on *Celosia argentea* (L.) with two plant essential oils. *Int J Pure Appl Zool* 2(3):187–195
- Enwezor WO, Udo EJ, Ayotade KA, Adepetu JA, Chude VO (1990) A review of soil and fertilizer use in Nigeria. In FPDD. Literature review on soil fertility investigations in Nigeria. (Volume 5 North West Zone). Federal Ministry of Agriculture and Natural Resources, Lagos, pp. 241–
- FMANR (Federal Ministry of Agriculture and Natural Resources) (1990) Literature review on soil fertility investigations in Nigeria. FMANR, Lagos
- Garbarino JR, Bednar AJ, Rutherford DW, Beyer RS, Wershaw RL (2003) Environmental fate of roxarsone in poultry litter. I. Degradation of roxarsone during composting. *Environ Sci Technol* 37:1509–1514. <https://doi.org/10.1021/es026219q>
- Hanč A, Tlustoš P, Száková J, Habart J, Gondek K (2008) Direct and subsequent effect of compost and poultry manure on the bioavailability of cadmium and copper and their uptake by oat biomass. *Plant Soil Environ* 54(7):271–278
- Jackson BP, Bertsch PM, Cabrera ML, Camberato JJ, Seaman JC, Wood CW (2003) Trace element speciation in poultry litter. *J Environ Qual* 32:535–540. <https://doi.org/10.2134/jeq2003.0535>
- Jayasuriya KMGG, Baskin J, Geneve RL, Baskin CC (2007) Morphology and anatomy of physical dormancy in *Ipomoea lacunosa*: identification of the water gap in seeds of Convolvulaceae (Solanales). *Ann Bot* 100:13–22. <https://doi.org/10.1093/aob/mcm070>
- Jonathan SG, Oyetunji OJ, Olawuyi OJ, Asemoloye MD (2012) Growth responses of *Corchorus olitorius* Lin. (Jute) to the application of organic manure as an organic fertilizer. *Acad Arena* 4(9):48–56
- Lindsay WI, Norvell WA (1978) Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Sc Soc Am J* 42:421–428. <https://doi.org/10.2136/sssaj1978.03615995004200030009x>
- Mazen A, Faheed FA, Ahmed AF (2010) Study of potential impacts of using organic manure in the amendment of desert reclaimed soil on wheat and Jews mallow plants. *Braz Arch Biol Technol* 53(4):917–930. <https://doi.org/10.1590/S1516-89132010000400022>
- Misra RV, Roy RN, Hiraoka H (2003) On-farm composting methods. Land and water discussion paper 2. Food and Agriculture Organization of the United Nations, Rome, pp 1–48
- Moral R, Paredes C, Bustamante MA, Marhuenda-Egea F, Bernal MP (2009) Utilisation of manure composts by high-value crops: safety and environmental challenges. *Bioresour Technol* 100(22):5454–5460. <https://doi.org/10.1016/j.biortech.2008.12.007>
- Nelson DW, Sommer LE (1996) Total carbon, organic carbon and organic matter. In: Sparks DL (ed) Methods of soil analysis. Part 3. SSSA book series no 5. SSSA Madison, Wisconsin, pp 960–1010
- Ng CC, Rahman MM, Boyce AN, Abas (2016) Heavy metals phyto-assessment in commonly grown vegetables: water spinach (*I. aquatica*) and okra (*A. esculentus*). *SpringerPlus* 5:469. <https://doi.org/10.1186/s40064-016-2125-5>
- Ogutade OA, Adebare TO, Odusanya OA, Oduntan OB (2015) The role of composted poultry manure in reducing bioavailability of heavy metals in tissues of *Celosia argentea* L. grown on dumpsite and garden soils. In: Rahmann G, Olabiyi TI, Olowe VIO (eds) Scientific track proceedings of the third african organic conference, Lagos, Nigeria, pp 89–92
- Okalebo JR, Gathua KW, Woomer PL (1993) Laboratory methods of soil and plant analysis: a working manual. TSBF, Nairobi
- Oyedele DJ, Asonugbo C, Awotoye OO (2006) Heavy metals in soil and accumulated by edible vegetable after phosphate fertilizer application. *Electron J Environ Agric Food Chem* 5(4):1446–1453
- Petersen SO, Henriksen K, Mortensen GK, Krogh PH, Brandt KK, Sorensen J, Madsen T, Petersen J, Gron C (2003) Recycling of sewage sludge and household compost to arable land: fate and effects of organic contaminants, and impact on soil fertility. *Soil Tillage Res* 72:139–152. [https://doi.org/10.1016/S0167-1987\(03\)00084-9](https://doi.org/10.1016/S0167-1987(03)00084-9)
- Samra I, Piliz S, Ferdag C (2007) Antibacterial and antifungal activity of *Corchorus olitorius* L. (Molokhia) extracts. *Int J Nat Eng Sci* 1(3):39–61
- Sapkota AR, Lefferts LY, McKenzie S, Walker P (2007) What do we feed to food-production animals? A review of animal feed ingredients and their potential impacts on human health. *Environ Health Perspect* 115:663–670. <https://doi.org/10.1289/ehp.9760>
- Schippers RR (2002) African indigenous vegetables. An overview of the cultivated species. Natural Resources International Limited and 94 Horticultural Development Services. Natural Resources Institute/ACP-EU Technical Centre for Agricultural and Rural Cooperation, Chatham
- Sims JT, Johnson GV (1991) Micronutrient soil test. In: Mortvedt JJ, Cox FR, Welch LM (eds) Micronutrients in agriculture, vol 4, 2nd edn. Soil Science Society of America book series. Soil Science Society of America, Madison
- Sobulo RA, Osiname OA (1981) Soils and fertilizer use in Western Nigeria. *Res Bull IART Ib* 11:20–26
- Tlustoš P, Szakova J, Korinek K, Pavlikova D, Hanc A, Balik J (2006) The effect of liming on cadmium, lead and zinc uptake reduction by spring wheat grown in contaminated soil. *Plant Soil Environ* 52:16–24
- Tolorunse KD, Ibrahim H, Aliyu NC, Oladiran JA (2015) The quality of jute mallow seeds exposed to different hot water-steeping and cooling protocols. *Am J Exp Agric* 7(2):107–117. <https://doi.org/10.9734/AJEA/2015/12517>
- Udo EJ, Ibis TO, Ogunwale JA, Ano AO, Esu IE (2009) Manual of Soil, Plant and Water Analysis. Sibon Books Ltd, Lagos
- Udoh BT, Ibia TO, Udo BU, Edem SO (2008) Assessment of micronutrient status of inland depression and floodplain (wetland) soils

- in Akwa Ibom State, South eastern Nigeria. *J Trop Agric Food Environ Ext* 7(2):156–161
- Uponi JJ, Adeoye GO (2000) Soil testing and plant analysis: an overview. In: Akoroda MO (ed) *Agronomy in Nigeria*. Department of Agronomy, University of Ibadan, Ibadan, pp 174–182
- van Zanten HHE, Mollenhorst H, de Vries JW, van Middelaar CE, van Kernebeek HRJ, de Boer IJM (2014) Assessing environmental consequences of using co-products in animal feed. *Int J Life Cycle Assess* 19:79–88. <https://doi.org/10.1007/s11367-013-0633-x>
- Walkley A, Black CA (1934) An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci* 37:29–38. <https://doi.org/10.1097/00010694-193401000-00003>
- Wang XH, Song YP, Ren J, Liu TK, Hou XL, Li Y (2013) Response of biomass and photosynthesis in non-heading Chinese cabbage excess copper. *J Anim Plant Sci* 23:1659–1665
- Yang XE, Long XX, Ni WZ, Ye ZQ, He ZL, Stoffella PJ, Calvert DV (2002) Assessing copper thresholds for phytotoxicity and potential dietary toxicity in selected vegetable crops. *J Environ Sci Health B* 37:625–635. <https://doi.org/10.1081/PFC-120015443>
- Zhang SQ, Zhang FD, Liu XM, Wang YJ, Zou SW, He XS (2005) Determination and analysis on main harmful composition in excrement of scale livestock and poultry feedlots. *Plant Nutr Fert Sci* 11:822–829

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

